

Non-singlet Q-deformations of $\mathcal{N} = 2$ gauge theories *

A. De Castro ^a, E. Ivanov ^b, O. Lechtenfeld ^a and L. Quevedo ^a

^a Institut für Theoretische Physik, Universität Hannover,
 Appelstrasse 2, 30167 Hannover, Germany.
 castro, lechtenf, quevedo@itp.uni-hannover.de

^b Bogoliubov Laboratory of Theoretical Physics, JINR,
 141980 Dubna, Moscow Region, Russia.
 eivanov@theor.jinr.ru

Abstract

We study a non-anticommutative chiral non-singlet deformation of the $\mathcal{N}=(1,1)$ abelian gauge multiplet in Euclidean harmonic superspace. We present a closed form of the gauge transformations and the unbroken $\mathcal{N}=(1,0)$ supersymmetry transformations preserving the Wess-Zumino gauge, as well as the bosonic sector of the $\mathcal{N}=(1,0)$ invariant action. This contribution is a summary of our main results in hep-th/0510013.

Extensions of gauge theories to non-commutative and non-(anti)commutative superspaces are currently of remarkable interest within the high energy physics community, mainly due to their relevance to subjects like string theory (see for example [2, 3] and references therein). Here we focus in a subclass of non-(anti)commutative Euclidean supersymmetric field theories called *Q-deformed*, realized via a Weyl-Moyal product with a bilinear nilpotent Poisson operator, which is constructed in terms of the supercharges,

$$P = -\overleftarrow{Q}_\alpha^i C_{ik}^{\alpha\beta} \overrightarrow{Q}_\beta^k. \quad (1)$$

The Moyal product of two superfields is then defined by

$$A \star B = Ae^P B. \quad (2)$$

The deformation parameters $C_{ij}^{\alpha\beta}$ form a constant tensor which is symmetric under the simultaneous permutation of the Latin and Greek indices, $C_{ij}^{\alpha\beta} = C_{ji}^{\beta\alpha}$. Generically, it breaks the full automorphism symmetry $Spin(4) \times O(1,1) \times SU(2) \equiv SU(2)_L \times SU(2)_R \times O(1,1) \times SU(2)$ of the $\mathcal{N} = (1,1)$ superalgebra — $O(1,1)$ and $SU(2)$ being the R-symmetry groups — down to $SU(2)_R$. An important feature of Q-deformations is the nilpotent nature

*To be published in the *Proceedings of the International Workshop "Supersymmetries and Quantum Symmetries"*, JINR, Dubna, 2005

of the Poisson operator ($P^5 = 0$) which makes the Moyal product polynomial, ensuring local actions. In virtue of the commutation properties of P with respect to the spinor covariant derivatives,

$$[D_\alpha^\pm, P] = 0, \quad [\bar{D}_\alpha^\pm, P] = 0, \quad [D^{\pm\pm}, P] = 0, \quad (3)$$

the product (2) breaks $\mathcal{N} = (1, 1)$ supersymmetry down to $\mathcal{N} = (1, 0)$ while preserving both chirality and Grassmann harmonic analyticity of the involved superfields, as well as the harmonic conditions¹ $D^{\pm\pm}A = 0$. Operator (1) can be split as follows,

$$P = -I \overleftarrow{Q}_\alpha^i \varepsilon^{\alpha\beta} \varepsilon_{ik} \overrightarrow{Q}_\beta^k - \overleftarrow{Q}_\alpha^i \hat{C}_{ik}^{\alpha\beta} \overrightarrow{Q}_\beta^k. \quad (4)$$

The first term is $Spin(4) \times SU(2)$ -preserving while the second term involves a $SU(2)_L \times SU(2)$ constant tensor which is symmetric under the independent permutations of Latin and Greek indices, $\hat{C}_{ij}^{\alpha\beta} = \hat{C}_{ij}^{\beta\alpha} = \hat{C}_{ji}^{\alpha\beta}$. For the generic choice, it fully breaks Euclidean symmetry, $SU(2)_L$ and R-symmetry $SU(2)$. Q-deformations induced only by the first term are called *singlet* or *QS-deformations*, whereas those associated with the second term, *non-singlet* or *QNS-deformations*. In this contribution we report important results on dynamical aspects of *QNS-deformations* of the $N = (1, 1)$ $U(1)$ vector multiplet in harmonic superspace. The talk is based on paper [1], where detailed calculations are performed and a complete list of references is given.

Gauge transformations. The residual gauge transformations of the component fields of the Abelian $\mathcal{N} = (1, 1)$ vector multiplet in the WZ gauge can be found from the Q-deformed superfield transformation [8]

$$\delta_\Lambda V_{WZ}^{++} = D^{++}\Lambda + [V_{WZ}^{++}, \Lambda]_\star, \quad (5)$$

with V_{WZ}^{++} being the analytic harmonic $U(1)$ superfield gauge connection and Λ the analytic residual gauge parameter satisfying $D_\alpha^+\Lambda = \bar{D}_\alpha^+\Lambda = 0$. In the left-chiral basis, where $x_A^{\alpha\dot{\alpha}} = x_L^{\alpha\dot{\alpha}} - 4i\theta^{-\alpha}\bar{\theta}^{+\dot{\alpha}}$ [9], V_{WZ}^{++} has the following θ -expansion

$$\begin{aligned} V_{WZ}^{++} = & (\theta^+)^2 \bar{\phi} + \bar{\theta}_\alpha^+ [2\theta^{+\alpha} A_\alpha^{\dot{\alpha}} + 4(\theta^+)^2 \bar{\Psi}^{-\dot{\alpha}} - 2i(\theta^+)^2 \theta^{-\alpha} \partial_\alpha^{\dot{\alpha}} \bar{\phi}] \\ & + (\bar{\theta}^+)^2 \left[\phi + 4\theta^+ \Psi^- + 3(\theta^+)^2 D^{--} - i(\theta^+ \theta^-) \partial^{\alpha\dot{\alpha}} A_{\alpha\dot{\alpha}} + \theta^{-\alpha} \theta^{+\beta} F_{\alpha\beta} \right. \\ & \left. - (\theta^+)^2 (\theta^-)^2 \square \bar{\phi} + 4i(\theta^+)^2 \theta^{-\alpha} \partial_{\alpha\dot{\alpha}} \bar{\Psi}^{-\dot{\alpha}} \right]. \end{aligned} \quad (6)$$

The superparameter $\Lambda_0 = ia + 2\theta^{-\alpha}\bar{\theta}^{+\dot{\alpha}}\partial_{\alpha\dot{\alpha}}a - i(\theta^-)^2(\bar{\theta}^+)^2\square a$ (being a an arbitrary function of x_L) found for the undeformed and singlet cases, breaks the WZ gauge in the non-singlet case, due to the appearance of an unwanted dependence on the harmonic variables $u_i^\pm$ in the expressions for the gauge variations. It is clear that is imperative to choose a gauge parameter Λ that preserves the WZ gauge, that is, for non-singlet deformations, some correction terms $\Delta\Lambda$ must be added to Λ_0 , where

$$\begin{aligned} \Delta\Lambda = & \theta_\alpha^+ \bar{\theta}_\alpha^+ \partial_\beta^{\dot{\alpha}} a B_1^{-\alpha\beta} + (\bar{\theta}^+)^2 \partial_{\beta\dot{\beta}} a A_\alpha^{\dot{\beta}} G^{--\alpha\beta} + (\theta^+)^2 (\bar{\theta}^+)^2 \square a P^{-4} \\ & + (\bar{\theta}^+)^2 \theta_\alpha^+ \left[\bar{\Psi}^{-\dot{\beta}} \partial_{\beta\dot{\beta}} a B_2^{-\alpha\beta} + \bar{\Psi}^{+\dot{\beta}} \partial_{\beta\dot{\beta}} a G^{-4\alpha\beta} \right] + (\theta^+)^2 (\bar{\theta}^+)^2 \partial_{\alpha\dot{\alpha}} a \partial_\beta^{\dot{\alpha}} \bar{\phi} B_3^{-4\alpha\beta} \\ & + i\theta_\alpha^+ \theta_\beta^- (\bar{\theta}^+)^2 \square a B_1^{-\alpha\beta} + i\theta_\alpha^+ \theta_\gamma^- (\bar{\theta}^+)^2 \partial_{\beta\dot{\lambda}} a \partial^{\gamma\dot{\lambda}} \bar{\phi} \frac{d}{d\phi} B_1^{-\alpha\beta}. \end{aligned} \quad (7)$$

¹Giving up chirality and analyticity, it is also possible to use spinor covariant derivatives to construct a nilpotent Poisson operator. We recommend ref.[5] for a deeper treatment of the subject.

The coefficients in (7) are some undetermined functions of harmonics, the field $\bar{\phi}$ and deformation parameters, calculated by requiring

$$\partial^{++}\delta A_{\alpha\dot{\alpha}} = 0, \quad \partial^{++}\delta\phi = 0, \quad (\partial^{++})^2\delta\Psi_{\alpha}^{-} = 0, \quad (\partial^{++})^3\delta D^{--} = 0. \quad (8)$$

The correction term to $\delta_0 V_{\text{WZ}}^{++}$ is,

$$\hat{\delta} V_{\text{WZ}}^{++} = D^{++}\Delta\Lambda + [V_{\text{WZ}}^{++}, \Delta\Lambda]_{\star}, \quad (9)$$

and the full WZ preserving gauge transformations are given by $\delta V_{\text{WZ}}^{++} = \delta_0 V_{\text{WZ}}^{++} + \hat{\delta} V_{\text{WZ}}^{++}$. Unfortunately, it is very difficult to find closed solutions of these equations for general deformation parameters, though their perturbative solutions always exist as series expansion. For the general choice of $\hat{C}_{\alpha\beta}^{ij}$, the gauge and susy transformations and the corresponding action are known only to few first orders in the parameters of deformation [6]. Nevertheless, exact solutions can be found for the product structure

$$\hat{C}_{\alpha\beta}^{ij} = b^{ij} c_{\alpha\beta},$$

which corresponds to the maximally symmetric non-singlet deformations. The full set of non trivial QNS-deformed gauge transformations laws for the $\mathcal{N} = (1, 1)$ vector multiplet in WZ gauge are then

$$\delta\bar{\phi} = 0, \quad \delta\bar{\Psi}_{\dot{\alpha}}^k = 0, \quad (10)$$

$$\delta A_{\alpha\dot{\alpha}} = X \coth X \partial_{\alpha\dot{\alpha}} a, \quad (11)$$

$$\delta\phi = 2\sqrt{c^2 b^2} \left(\frac{1 - X \coth X}{X} \right) A^{\alpha\dot{\alpha}} \partial_{\alpha\dot{\alpha}} a, \quad (12)$$

$$\begin{aligned} \delta\Psi_{\alpha}^i = & \left\{ \left[\frac{4X^2(X \coth X - 1)}{X^2 + \sinh^2 X - X \sinh 2X} \right] b^{ij} c_{\alpha\beta} \right. \\ & \left. - \sqrt{c^2 b^2} \left[\frac{4X \cosh^2 X - 2X^2(\coth X + X) - \sinh 2X}{X^2 + \sinh^2 X - X \sinh 2X} \right] \varepsilon^{ij} \varepsilon_{\alpha\beta} \right\} \bar{\Psi}_{j\dot{\alpha}} \partial^{\beta\dot{\alpha}} a. \end{aligned} \quad (13)$$

$$\delta D_{ij} = 2ib_{ij} c^{\alpha\beta} \partial_{\alpha\dot{\alpha}} \bar{\phi} \partial_{\beta}^{\dot{\alpha}} a. \quad (14)$$

where

$$X = 2\bar{\phi} \sqrt{b^{ij} b_{ij} c_{\alpha\beta}^{\alpha\beta}}. \quad (15)$$

Detailed calculations of these transformations laws are carried out in [1]. Having the explicit QNS-deformed gauge transformations, one can deduce a minimal Seiberg-Witten-like map which puts these transformations into the standard undeformed form

$$\begin{aligned} \Psi_{\alpha}^i &= \tilde{\Psi}_{\alpha}^i + 2\sqrt{c^2 b^2} \left[2 \left(\coth X - \frac{1}{X} \right) - X \right] \bar{\Psi}^{i\dot{\alpha}} \tilde{A}_{\alpha\dot{\alpha}}, \quad D_{ij} = \tilde{D}_{ij} + 2ib_{ij} c^{\alpha\beta} \partial_{\alpha\dot{\alpha}} \bar{\phi} \tilde{A}_{\beta}^{\dot{\alpha}}, \\ A_{\alpha\dot{\alpha}} &= \tilde{A}_{\alpha\dot{\alpha}} X \coth X, \quad \phi = \tilde{\phi} + \tilde{A}^2 \sqrt{c^2 b^2} X \coth X \left(\frac{1 - X \coth X}{X} \right). \end{aligned} \quad (16)$$

For the fields with tilde we obtain the standard transformations

$$\delta\tilde{A}_{\alpha\dot{\alpha}} = \partial_{\alpha\dot{\alpha}} a, \quad \delta\tilde{\phi} = 0, \quad \delta\tilde{D}^{ij} = 0, \quad \delta\tilde{\Psi}_{\alpha}^k = 0.$$

The gauge field strength $F_{\alpha\beta} = 2i\partial_{(\alpha\dot{\alpha}}\dot{A}_{\beta)}^{\dot{\alpha}}$ which is non-covariant with respect to the deformed transformations is redefined under the transformation $A_{\alpha\dot{\alpha}} \rightarrow \tilde{A}_{\alpha\dot{\alpha}}$ as

$$F_{\alpha\beta} = \tilde{F}_{\alpha\beta} X \coth X + 4i\sqrt{b^2 c^2} \tilde{A}_{(\beta\dot{\alpha}} \partial_{\alpha)}^{\dot{\alpha}} \bar{\phi} \left(\coth X - \frac{X}{\sinh^2 X} \right), \quad \text{where} \quad \tilde{F}_{\alpha\beta} = 2i\partial_{(\alpha\dot{\alpha}} \tilde{A}_{\beta)}^{\dot{\alpha}}.$$

Unbroken susy transformations. Unbroken supersymmetry is realized on V_{WZ}^{++} as

$$\delta V_{\text{WZ}}^{++} = (\epsilon^{+\alpha} \partial_{+\alpha} + \epsilon^{-\alpha} \partial_{-\alpha}) V_{\text{WZ}}^{++} - D^{++} \Lambda_c - [V_{\text{WZ}}^{++}, \Lambda_c]_{\star}, \quad (17)$$

where the star bracket, like in the previous consideration, is defined via the non-singlet Poisson structure with the deformation matrix $\hat{C}_{\alpha\beta}^{ij} = b^{ij} c_{\alpha\beta}$ and Λ_c is the compensating gauge parameter which is necessary for preserving WZ gauge. As in the QNS-deformed gauge transformations case, variations obtained using the original $\mathbf{L}$ for undeformed and singlet cases (see[8]), violate the WZ gauge due to an unbalanced appearance of harmonic and Grassmann variables [1], so one is led to properly modify $\mathbf{L}$. Thus we define

$$\Lambda_c = \mathbf{L} + F_{\epsilon} \quad (18)$$

We denote by $\delta V_{\text{WZ}}^{++}$ the lowest-order non-singlet part of the transformations coming from the star commutator in (17) using $\mathbf{L}$, and rewrite (17) in the following way

$$\delta V_{\text{WZ}}^{++} = \delta V_{\text{WZ}}^{++} - D^{++} F_{\epsilon} - [V_{\text{WZ}}^{++}, F_{\epsilon}]_{\star} \quad (19)$$

with

$$\begin{aligned} \mathbf{L} = & 2(\epsilon^{-}\theta^{+})\bar{\phi} + \bar{\theta}_{\dot{\alpha}}^{-} [4i(\epsilon^{-}\theta^{+})\theta_{\alpha}^{-}\partial^{\alpha\dot{\alpha}}\bar{\phi} - 2\epsilon_{\alpha}^{-}A^{\alpha\dot{\alpha}} + 4(\epsilon^{-}\theta^{+})\bar{\Psi}^{-\dot{\alpha}}] + \\ & (\bar{\theta}^{+})^2 \left[2(\epsilon^{-}\Psi^{-}) + 2i\epsilon^{-\alpha}\theta^{-\beta}\partial_{\beta}^{\dot{\alpha}}A_{\alpha\dot{\alpha}} - 2(\epsilon^{-}\theta^{+})(\theta^{-})^2\Box\bar{\phi} \right. \\ & \left. + 4i(\epsilon^{-}\theta^{+})\theta^{-\alpha}\partial_{\alpha\dot{\alpha}}\bar{\Psi}^{-\dot{\alpha}} + 2(\epsilon^{-}\theta^{+})D^{--} \right]. \end{aligned} \quad (20)$$

The additional compensating gauge parameter intended for restoring the WZ gauge with the minimal set of terms needed to eliminate the improper harmonic and Grassmann dependence amounts to the following form

$$\begin{aligned} F_{\epsilon} = & \theta^{+\alpha} f_{\alpha}^{-} + \bar{\theta}_{\dot{\alpha}}^{+} [\bar{g}^{-\dot{\alpha}} + 2i\theta_{\alpha}^{-}\theta^{+\beta}\partial^{\alpha\dot{\alpha}} f_{\beta}^{-} + \theta^{+\alpha} b_{\alpha}^{-\dot{\alpha}} + (\theta^{+})^2 \bar{g}^{(-3)\dot{\alpha}}] \\ & + (\bar{\theta}^{+})^2 \left[g^{--} - (\theta^{-})^2 \theta^{+\alpha} \Box f_{\alpha}^{-} + i\theta^{-\alpha} \partial_{\alpha\dot{\alpha}} \bar{g}^{-\dot{\alpha}} + i\theta^{+\alpha} \theta^{-\beta} \partial_{\beta}^{\dot{\alpha}} b_{\alpha\dot{\alpha}}^{-} + \theta^{+\alpha} f_{\alpha}^{(-3)} \right. \\ & \left. + i(\theta^{+})^2 \theta^{-\alpha} \partial_{\alpha\dot{\alpha}} \bar{g}^{(-3)\dot{\alpha}} + (\theta^{+})^2 X^{(-4)} \right]. \end{aligned} \quad (21)$$

Requiring the elimination of terms with unbalanced Grassmann variables and

$$\partial^{++}\delta\bar{\phi} = 0, \quad (\partial^{++})^2 \delta\bar{\Psi}_{\dot{\alpha}}^{-} = 0, \quad \partial^{++}\delta A_{\alpha\dot{\alpha}} = 0, \quad (\partial^{++})^2 \delta\Psi_{\alpha}^{-} = 0, \quad (\partial^{++})^3 \delta D^{--} = 0, \quad (22)$$

we can explicitly find components of F_{ϵ} and restore the correct $\mathcal{N} = (1, 0)$ supersymmetry transformations preserving WZ gauge. The full set of these transformations together with the full supersymmetric action will be given² in [10]. Here we show the simplest subalgebra

$$\begin{aligned} \delta\bar{\phi} = & 0, \quad \delta A_{\alpha\dot{\alpha}} = 8\bar{\phi}\epsilon^{i\beta}\bar{\Psi}_{\dot{\alpha}}^j b_{ij} c_{\alpha\beta} + 2\epsilon_{\alpha}^k \bar{\Psi}_{k\dot{\alpha}} X \coth X, \\ \delta\bar{\Psi}_{\dot{\alpha}}^i = & \left[\frac{2i}{\sqrt{b^2 c^2}} \cosh X \sinh X c^{\alpha\beta} b^{ij} - i \cosh^2 X \varepsilon^{\alpha\beta} \varepsilon^{ij} \right] \epsilon_{j\beta} \partial_{\alpha\dot{\alpha}} \bar{\phi}. \end{aligned} \quad (23)$$

²In fact, it is of no actual necessity to explicitly know these transformations, since our procedure of deriving the action is manifestly $\mathcal{N} = (1, 0)$ supersymmetric by construction [1].

These variations form an algebra which is closed modulo a gauge transformation with the composite parameter $a_c = -2i(\epsilon \cdot \eta)\bar{\phi}$:

$$[\delta_\epsilon, \delta_\eta]\bar{\phi} = 0, \quad [\delta_\epsilon, \delta_\eta]\bar{\Psi}_\alpha^j = 0, \quad [\delta_\epsilon, \delta_\eta]A_{\alpha\dot{\alpha}} = -2i(\epsilon \cdot \eta)(X \coth X)\partial_{\alpha\dot{\alpha}}\bar{\phi}.$$

Bosonic action. Now we present the bosonic sector of the $\mathcal{N} = (1, 0)$ gauge invariant action in components. The QNS-deformed action for the $\mathcal{N} = (1, 1)$ U(1) gauge theory in harmonic superspace [9], in the form most appropriate for our purposes, is written in the same way as in the QS-deformed case [8]

$$S = \frac{1}{4} \int d^4x_L d^4\theta du \mathcal{W} \star \mathcal{W} = \frac{1}{4} \int d^4x d^4\theta du \mathcal{W}^2. \quad (24)$$

Here $\mathcal{W}$ is the covariant superfield strength

$$\mathcal{W} = -\frac{1}{4}(\bar{D}^+)^2 V^{--} \equiv \mathcal{A}(x_L, \theta^+, \theta^-) + \bar{\theta}_\alpha^+ \tau^{-\dot{\alpha}}(x_L, \theta^+, \theta^-) + (\bar{\theta}^+)^2 \tau^{--}(x_L, \theta^+, \theta^-), \quad (25)$$

and V^{--} is the non-analytic harmonic connection related to V_{WZ}^{++} by the harmonic flatness condition

$$D^{++}V^{--} - D^{--}V_{\text{WZ}}^{++} + [V_{\text{WZ}}^{++}, V^{--}]_\star = 0. \quad (26)$$

The whole effect of the considered deformation in the above action comes from the structure of $\mathcal{W}$ due to the presence of the star commutator in the equation (26) defining V^{--} . As a consequence of the latter, (25) satisfies the condition

$$D^{++}\mathcal{W} + [V_{\text{WZ}}^{++}, \mathcal{W}]_\star = 0. \quad (27)$$

It is not hard to prove that the only contribution to the entire action is the superfield $\mathcal{A}$ in (25) (see [8]). Thus, the invariant action is reduced to

$$S = \frac{1}{4} \int d^4x d^4\theta du \mathcal{A}^2. \quad (28)$$

Once again we refer to [1] for details of the calculations leading to the relevant components of $\mathcal{A}$. Finally the bosonic limit of the action, after performing the minimal SW map (16), is

$$S_{\text{bos}} = \int d^4x \left[-\frac{1}{2}\tilde{\phi}\square\bar{\phi} - \frac{1}{2}(b^2 c^2)^{3/2} \tanh X \partial_{\alpha\dot{\alpha}}\bar{\phi} \partial^{\alpha\dot{\alpha}}\bar{\phi} \square\bar{\phi} + \frac{1}{4} \frac{\tilde{D}^2}{\cosh^2 X} - \frac{1}{16} \tilde{F}^2 \cosh^2 X + \frac{1}{4} b^2 (c \cdot \tilde{F})^2 \bar{\phi}^2 \frac{\sinh^2 X}{X^2} + \frac{1}{2} \bar{\phi} (b \cdot \tilde{D}) (c \cdot \tilde{F}) \frac{\tanh X}{X} \right]. \quad (29)$$

This action is invariant under the standard abelian gauge transformations. Turning off the deformation parameters we are left with the usual bosonic sector of the undeformed action. Performing the further field redefinition

$$d^{ij} = \frac{1}{\cosh^2 X} \tilde{D}^{ij} + \bar{\phi} (c \cdot \tilde{F}) b^{ij} \frac{\tanh X}{X}, \quad \varphi = \frac{1}{\cosh^2 X} \left[\tilde{\phi} + (b^2 c^2)^{3/2} (\partial\bar{\phi})^2 \tanh X \right],$$

the bosonic action can be transformed into a simple form

$$S_{bos} = \int d^4x \cosh^2 X \left[-\frac{1}{2} \varphi \square \bar{\phi} + \frac{1}{4} d^{ij} d_{ij} - \frac{1}{16} \tilde{F}^{\alpha\beta} \tilde{F}_{\alpha\beta} \right]. \quad (30)$$

From this expression it is obvious that we cannot disentangle the interaction between the gauge field and $\bar{\phi}$ by any field redefinition. This is similar to the singlet case [8, 7], where a scalar factor $(1 + 4I\bar{\phi})^2$ appears instead of $\cosh^2 X$. Note that the bosonic action (30) involves only squares c^2 and b^2 , so it preserves space-time $Spin(4) = SU(2)_L \times SU(2)_R$ symmetry and $SU(2)$ R-symmetry as in the singlet case. This property is similar to what happens in the deformed Euclidean $\mathcal{N} = (1/2, 1/2)$ Wess-Zumino model where the deformation parameter $C^{\alpha\beta}$ also appears squared [3]. However, we know that the fermionic completion of (30) will explicitly include both $c^{\alpha\beta}$ and b^{ik} [10], so these two symmetries are broken in the total action. This feature also compares with the breaking of Lorentz symmetry in the deformed $\mathcal{N} = (1/2, 1/2)$ gauge theory action, due to fermionic terms [3]. Though the string interpretation of the QS-deformation is known [8], the possible stringy origin of the non-singlet case —e.g. as some special $\mathcal{N} = 4$ superstring background— is still unclear.

References

- [1] A. De Castro, E. Ivanov, O. Lechtenfeld and L. Quevedo, [arXiv:hep-th/0510013].
- [2] M. R. Douglas and N. A. Nekrasov, Rev. Mod. Phys. **73** (2001) 977 [arXiv:hep-th/0106048]; J. de Boer, P. A. Grassi and P. van Nieuwenhuizen, Phys. Lett. **B 574** (2003) 98 [arXiv:hep-th/0302078]; H. Ooguri and C. Vafa, Adv. Theor. Math. Phys. **7** (2003) 53 [arXiv:hep-th/0302109]; Adv. Theor. Math. Phys. **7** (2004) 405 [arXiv:hep-th/0303063];
- [3] N. Seiberg, JHEP **0306** (2003) 010 [arXiv:hep-th/0305248].
- [4] E. Ivanov, O. Lechtenfeld and B. Zupnik, JHEP **0402** (2004) 012 [arXiv:hep-th/0308012].
- [5] L. Tamassia, “Noncommutative supersymmetric / integrable models and string theory,” PhD thesis, [arXiv:hep-th/0506064]; S.V. Ketov and S. Sasaki, Phys. Lett. **B 595** (2004) 530 [ArXiv:hep-th/0404119]; Phys. Lett. **B 597** (2004) 105 [ArXiv:hep-th/0405278]; Int. J. Mod. Phys. **A 20** (2005) 4021 [arXiv:hep-th/0407211]; S. Ferrara, M.A. Lledo and O. Macia, JHEP **0309** (2003) 068 [arXiv:hep-th/0307039].
- [6] T. Araki and K. Ito, Phys. Lett. **B 595** (2004) 513 [arXiv:hep-th/0404250]; T. Araki, K. Ito and A. Ohtsuka, JHEP **0401** (2004) 046 [arXiv:hep-th/0401012]; T. Araki, K. Ito and A. Ohtsuka, Phys. Lett. **B 606** (2005) 202 [arXiv:hep-th/0410203]; T. Araki, K. Ito and A. Ohtsuka, JHEP **0505** (2005) 074 [arXiv:hep-th/0503224].
- [7] T. Araki, K. Ito and A. Ohtsuka, JHEP **0401** (2004) 046 [arXiv:hep-th/0401012].
- [8] S. Ferrara, E. Ivanov, O. Lechtenfeld, E. Sokatchev and B. Zupnik, Nucl. Phys. **B 704** (2005) 154 [arXiv:hep-th/0405049].

- [9] A.S. Galperin, E.A. Ivanov, V.I. Ogievetsky and E.S. Sokatchev, *Harmonic Superspace*, Cambridge University Press 2001, 306 p.
- [10] A. De Castro, E. Ivanov, O. Lechtenfeld and L. Quevedo, work in preparation.
- [11] T. Araki, T. Takashima and S. Watamura, arXiv:hep-th/0510088.